

No. 64

No. 64

Hydrocephalus
12

Hydrocephalus
Hydrothorax

James W. M. Wallace

admitted March 23. 1819.

Ms. A. 9. 11

James H. McMillan
Secretary of the Board of Trustees

It is
of a subject
choose one
hope to come
and that are
ment of the
this view, he
by offering a
cases and ca
of nature of
enlarged co
which better
ment and a pr
rejection I do
of physics. N
by underscri
candidate for
ation in which
pleasure on the
worked in diff
light on the

Observations on Hydrothorax.

It can scarcely be expected that in selecting a subject for an inaugural dissertation I should choose one in the investigation of which I could hope to correct former error, or advance any thing new that would have for its object the improvement of the science of medicine. Attempts with this view, however desirable, I at present decline by offering some general observations on the seat, causes and cure of hydrothorax; leaving to those of maturer years, sounder judgment and more enlarged experience to explain the difficulties which hitherto have prevented the full development and perfection of "the healing art." By this expression I do not mean to deprecate the profession of physic. The very reverse would be my wish. By embracing in it, and presenting myself as a candidate for a degree, I sufficiently show the estimation in which I hold it; whilst I dwell with pleasure on the contributions that have been furnished in different ages and with no little delight on the improvement which medicine has

received
from the
being a
Before
short of de
law of the
induced m
a vector; d
inexpensive
guarantee
of allowance
I collect
the cavity of
question-w
in one or
inexpensive
provisional
definition
subject, but
purpose of
varied as
tion of the
not be adop
Hydrothorax
was; yet in

received in this country, and in an especial manner from the University in which I have the honour of being a student.

Before I proceed further, I must take the liberty of stating, that nothing, short of a positive law of the University requiring it, could ever have induced me to appear before the bust of judges as a writer, disqualified as I feel myself by age and inexperience. - Then, I indulge the hope, well guaranteed to this performer, the full share of allowance its imperfections demand.

A collection of serous fluid in any part of the cavity of the thorax constitutes the disease in question - whether the fluid effused be present in one or both cavities of the pleura, in the parenchymatous structure of the lung, or in the pericardium alone. I am not certain that this definition is sanctioned by the writers on this subject, but as it appears to me to answer all the purposes of practice, (the treatment not being varied so far as I know by the different situation of the water) I see no reason why it should not be adopted.

Hydrothorax is not a disease of very frequent occurrence; yet in every point of view it merits the par-

[Faint, mostly illegible handwriting in cursive script, likely bleed-through from the reverse side of the page.]

similar also
the form
of country
and may
I however
and arises
station. The
duration
am or by
but occasion
an opposit
usually we
attention
though, as
usually he
time, with
adjoining de
for the hum
recurrence
Symptoms
within the
self by con
tion of the
continued by
a experience

icular attention of the physician. It is usually, like other forms of dropsy, met with in marshy districts of country, succeeds intermittent and other fevers, and may be brought on from exposure to cold. It however, appears in all climates and situations, and arises in very different states of the constitution. Most frequently, at least as far as my observation has extended, it has been connected more or less with a phlogistic state of the system; but occasionally it is found to be present in an opposite condition of the constitution. Most usually we see this complaint blended with collections of water in other parts of the body; though unquestionably hydrothorax is occasionally known to exist for some length of time, without its being connected with, or inducing dropical effusions in any other part of the human structure. This however, is a rare occurrence.

Symptoms. An effusion of serous fluid within the thorax, usually first discovers itself by some uneasiness at the inferior portion of the sternum; which does not long continue before some difficulty of breathing is experienced. This is ordinarily for the first

[Faint, illegible handwriting on lined paper, likely bleed-through from the reverse side.]

time for
or when
during the
bold, the
constant
the com
is much
found,
plains of
of his son
say but
most. 7
dy of me
than the
most inf
of a flee
ing phys
the pater
in sudden
station of
With the
in common
intestinal
this adom
hyposole.

time felt, when the person ascends an acclivity, or when he places himself in a recumbent position. But after the disease has taken firm hold, the difficulty of breathing becomes more constant and very much annoys the patient. The urine is diminished in quantity. The thirst is much augmented. There is sometimes a fever present. The patient in some instances complains of a "lilative pain" or numbness in one of his arms. A cough comes on, that is at first dry, but after some continuance it becomes moist. The matter thrown off is composed chiefly of mucus. I think it is of a whiter colour than the saliva thrown off in health. The most infallible symptom is the fluctuation of a fluid, ascertained either by the attending physician or by the patient himself. The patient after going to sleep is subject to be suddenly awaked with starting and palpitation of the heart.

With the symptoms already enumerated there are commonly adematous swellings of the lower extremities. Some physicians have stated that this adema is often preceded by a partial hydrocele. The pulse, especially after the case

has been
and is inter-
posed
even at the
time. Mon-
is small and
uniformly
When
its progress
now and
the action
is by the
the fatal
suddenly to
instances of
crashed a
days before
Causes
one of hydro-
perhaps in
positive cer-
be expected
should be
long as the
in its power

has been of some standing) is small, irregular and intermitting. In one or two cases, that have presented themselves to my notice, the pulse even at this period was not without some tension. More frequently at this stage, the pulse is small and weak and the face is almost uniformly of a pale colour.

When ~~the~~ hydrothorax is not arrested in its progress, it commonly proceeds to give more and more difficulty of breathing, till the action of the lungs be entirely interrupted by the quantity of water effused; and the fatal event frequently happens more suddenly than was expected. In many of the instances of a fatal hydrothorax, I have remarked a spitting of blood to come on several days before the patient died. Cullen.

Causes. The causes of dropsy in general and of hydrothorax in particular are not perhaps in all cases to be pointed out with positive certainty; nor indeed ought it to be expected that our views on this head should be very clear and unequivocal, so long as the lymphatic system remains in its present unexplored state. We have,

A. I. Chapman inculcates a similar doctrine
in his lectures.

known,
with the e
direct us
clude now
fort will
cases can
sources of
Any co
tion, so tha
is implied
knowledge
as of disease
be mentio
blood-vessels
A. polyph a
impels conce
dication. A
known
and probably
causes operate
ticular ope
these causes
the production
of hydatids, a
near the head

however, in general, a sufficient acquaintance with the cause and nature of the disease to direct us successfully in its treatment. I allude now to some particular forms only, for it will be seen hereafter, that certain cases can not be expected to yield to the resources of our art.

Any cause operating on the venous system, so that the return of blood to the heart is impeded or prevented, is very uniformly acknowledged a cause of dropsy. Of the sources of disease operating in this manner, may be mentioned, spicification of the large blood-vessels or their valves; the existence of polypi and probably an atony of the vessels concerned in the parts under consideration.^A

Exposure to cold is to be considered as another, and probably not one of the least infrequent causes operating to the production of the particular species of dropsy before us. Of the other causes recognised as causing a power in the production of hydrothorax, are the presence of hydatids, aneurismal dilatations situated near the heart, and probably certain morbid

B. Maclean on Hydrothorax page 53.

C. I am fully aware an author^d of some distinction says, that he who will not admit the presence of a watery fluid in the chest without the fluctuation being heard will search for it in vain. The fluctuation of a fluid is certainly not always evident, but having seen some patients afflicted with hydrothorax who repeatedly told me that they both felt and heard the fluctuation in the chest in changes of posture, I can not doubt of its being a frequent symptom; and this opinion is sanctioned by the high authority of Cullen and others. *Lewros Cullens synopsis* p. 157.

B. Maclean p. 2nd.

conditions of that organ may excite the disease.

Dr Callen refers the existence of dropsy more immediately to "an increased secretion or to a diminished absorption". Perhaps both of these conditions sometimes exist.

Diagnosis. The complaint before us, is liable to be confounded with other diseases of the body.

Those affections for which it may be mistaken, are angina pectoris, asthma and aneurism of some of the large blood vessels of the heart. From the first of these hydrothorax may be distinguished by fluctuation of water in the breast; and by there not being connected with angina pectoris the edematous swellings of the lower extremities which we meet with when fluid has been effused in the thorax, especially if the disease be of some standing. There is also a diminution in the quantity of urine voided in dropsy a symptom I believe that never occurs in angina pectoris.

Hydrothorax may be confounded with asthma, but in general it can be distinguished by there not being a scarcity of urine in ~~the~~ asthma;

E. Cullens first lines Vol. 4th section 1702.

and
flo
sho
certain
sent,
and
breath
is d
now.

As
to the
be me
dotho
thorbe
of the
appears
the eve
and oc
case."

Harle
been the
Prog
has been
able. N
the aut

and by the paroxysm of this last named complaint pretty steadily coming on at night, whereas in dropsical effusions of the thorax certain symptoms are almost uniformly present, as paleness of the face, scarcity of urine, and in the last stage, the difficulty of breathing is always present. In asthma there is during the paroxysm a suffusion of face, never I believe present in hydrothorax.

Aneurism of the large arteries contiguous to the heart produces symptoms that may be mistaken for those which accompany hydrothorax. In general, however, a paul and throbbing in the seat and direction of some of the large arteries, never entirely ceasing, aggravated by the causes which accelerate the circulation of the blood, are experienced, and serve to distinguish it from this disease". Maclean p. 53.

Startings from sleep and palpitation have been thought to characterize hydrothorax.

Prognosis. In this affection the prognosis has been considered almost universally unfavourable. Were I to pronounce an opinion on the authority of authors, I should say a very

large for
would
own
clude
will be
consider
Could
state of
example
-pi in the
steris
to of
when it
was
individual
now in
- but
condition
remain
estimated
over a
terminat
In gen
judged
the

large proportion of the "unhappy sufferers" would not survive the disease. But from my own very limited observation, I should conclude that one half at least of all the cases will be found to yield to the influence of remedies.

Could we ascertain with precision on what state of the parts the disease depended, as for example when it owed its existence to polypus in the heart, to aneurisms of one of the large arteries immediately issuing from this organ, to ossification of these vessels or their valves, or when it came on as consequent to asthma, or was produced by hydatids we might with considerable certainty venture to predict the manner in which the complaint would terminate — but so long as the signs by which these conditions of the parts originating the disease remain in obscurity, so long must it be justly esteemed an exceedingly difficult task to deliver a correct opinion on what will be the termination of hydrothorax.

In general the degree of danger may be judged of from the difficulty of breathing, by the scarcity of urine, by the increasing excited

[Faint, mostly illegible handwriting in cursive script, spanning approximately 15 lines across the page.]

when the
stat poss
ers on ob
palpitatio
good de
ness.
We ha
the prog
the patie
ting drink
been brow
in advanc
mortal.
in practice
termination
been enu
a particula
infary.
Treatme
on the seat,
and probable
shall never
treatment, as
from the m
tions of the

when the patient puts his body in a horizontal position, by the difficulty he experiences in obtaining rest, by the violence of the palpitation, and above all by the little good derived from the exhibition of medicines.

We shall likewise be assisted in forming the prognosis, by ascertaining the habits of the patient in regard to the use of intoxicating drinks, and by his age. When it has been brought on by intemperance, and attacks in advanced life it almost certainly proves mortal. These are the circumstances by which, in practice, we are to direct our judgment on the termination of hydrothorax. Others might have been enumerated, but as they are more equivocal a particular notice of them is supposed unnecessary.

Treatment. Having made some observations on the seat, causes, distinguishing symptoms and probable termination of hydrothorax, we shall now proceed to the exposition of the treatment, which in a majority of cases, has proved the most salutary. All the investigations of the physician are directed to the

attainment of
not how long
have been so
in other depa
will avail be
of the proper
istence of de
In all ca
applied and co
they should,
do I have o
one prevail in
frame, so it w
will demand
ing as the on
do this does
atory nature
be required. In
to repeat this
morally it mo
reverts to it.
attack, in wh
collect eightie
help them twel
taken from th

attainment of this information; and it matters not how laboriously or successfully he may have been employed in obtaining correct views in other departments of his profession, the whole will avail but little if he remain ignorant of the proper treatment during the actual existence of disease.

In all cases where the cause can be distinguished and continue to operate on the body, they should, if practicable, be removed.

As I have said that hydrothorax may arise and prevail in two opposite states of the human frame, so it will be evident, these conditions will demand a different treatment according as the one or the other may prevail.

As this disease is most usually of an inflammatory nature, bleeding from the arm will be required. In common it is not necessary to repeat this evacuation often, but occasionally it may be necessary to have frequent recourse to it. I have myself witnessed an attack, in which there were as well I should collect eighteen bleedings promised, and not less than twelve or fourteen pounds of blood taken from the system. It is proper however

H. Derriars medical histories p. 347.

to show the
station,
case was a
complaint
or needs a
dies on last
found his
h. 1/2. This
before he was
us with ph
time opposi
operal sta
"Man 1/2
with the oap
tother" mea
breast and
insufficient st
H. case will
Chapmans
What to d
as I have ne
that much e
their applicat
They come
wor of the po

to observe here, that the patient was of large stature, had a full habit and that the disease was of many months continuance. The complaint during this period would advance or recede according as he employed the remedies or laid them aside. He very uniformly found his breathing rendered more free by V. S. This person ultimately recovered, but before he was reinstated to health, he was attacked with phlegmasia dolens a disease at one time supposed ^{to be} peculiar to women in the puerperal state.²

"When V. S. has been urged as far as consistent with the safety of the patient, we may resort to other means of depletion, as cups applied to the breast and sides. By this, alone, if used in the incipient stage of the disease, a great majority of cases will be arrested in their progress." Chapman's M. S. Lectures.

Next to bleeding blisters have been employed. I have never seen them used; but suppose that much benefit might be received from their application after the use of the lancet.

They come well recommended by the professor of the practice in the university of Penna.

G. Maclean p. 102.

-plethora.

Maclean

states that the

more speedily

to the observer

effects are two

Noting who

two are useful

much excitement

it might app

units admit of

to this disease

the necessary ge

it appears proba

be expected from

I shall hear to

Maclean's opinion

be between the

is separating to

tion of cups. They

are large enou

on some has be

in this way not

two blisters; and

=oxyhaemia.

Maclean in his treatise on hydrothorax, places blisters at the head of his remedies. He states that they relieve the urgent symptoms more speedily than the remedies addressed to the absorbents and kidneys, but says their effects are transient.

Acting upon the general principle that blisters are useless or inadmissible when there is much excitement of the sanguiferous system, it might appear doubtful whether they would admit of a very extensive application to this disease in its first stage; but after the necessary quantity of blood has been drawn, it appears probable that much benefit may be expected from the use of vesicatories.

I shall here take the liberty of copying Dr Chapman's opinion on this subject as delivered in his lectures. "Blisters" says he "may be regarded as cooperating to the same end with the application of cups. They should be repeatedly applied and large enough to cover the whole breast. As soon as one has healed, another should be applied. In this way nothing will be more efficacious than blisters; and to this point my own expe-

H. Maclean p. 117.

some forms
of
inflammation
are but of
which are the
cause in the
feels in this
and character
latent, it is
against ecto
rather loose
purpose are
the profusion
Notwithstanding
has varied and
time in height
coming of car
in this subject
former, one of
Myxocarpus, n
sinuifolia, b
which a mon
found in the
Bicolor.

Bicolor the

"since furnishes the most ample testimony."

Purgings, as being beneficial in other inflammatory affections, has been recommended here: but objections have been raised to its use, which are, that broides being useful or pernicious in the disorders of the chest generally, it fails in this instance of producing absorption, and exhausts too much the strength of the patient. It will be proper however, to guard against costiveness, and to keep the bowels rather loose than otherwise; and for this purpose cream of Tartar is probably to have the preference.

Notwithstanding the objections which have been urged against the employment of purgatives in Hydrothorax, I should perhaps, be deserving of censure were I to close the remarks on this subject, without observing that Dr Ferriar, one of the ablest of the British Physicians, not only thought purgatives admissible, but warmly ^{recommends} the Claterium, than which a more drastic cathartic is not to be found in the whole compass of the Materia Medica.

Besides the remedies already noticed, medicines

belonging to
each emp
nessy such
of their ad
monment, a
to each mon
that particu
We don't whe
ness of the
tous a day
on the orp
to acquire
ness in which
a close confid
a various com
abundant of
on two or three
view of the
of the patient
sions to each
on effects are o
shed by either
baptists have
which the ques
it contains

belonging to the class of diuretics have been much employed. They operate by increasing the urinary discharge.

Of these substances the squill perhaps stands preeminent. With a considerable diuretic power, it exerts more or less of an expectorant quality that particularly recommends it in this disease. The dose when taken alone, is from one to three grains of the dried root, given twice or three times a day in the form of pill. Notwithstanding the reputation which this medicine has acquired in dropsy, there are few if any cases in which ~~citrus-threo~~ or any other remedy is alone confided in. The squill is exhibited in various combinations. Combined with the submuriate of Mercury in the proportion of one, two or three grains of the former, to one grain of the latter, according to the state of the patient, and given twice a day it is found to exert its best powers. By such union effects are obtained that could not be produced by either singly.

Disputes have arisen as to the extent in which the squill is to be administered. By some it is contended that in order to ensure the

H. Chapman's Therapeutics & Mat. Med. Vol. 5th
p. 267.

diverted eff
own vomit
diverted ac
apportioned to
may proceed
ing the open
very tough l
the catenated
can will be
it ma
re action le

As respects
whole might
usage seems
number for
shape the w
brought in

The Lige
the list of
of the most
a cautious a
hardness eff
storming.
be best adap
four of artin

diuretic effect, it should produce nausea or even vomiting; whilst others assert, that the diuretic action neither depends on, nor is proportioned to the nausea or vomiting, which may proceed from its administration. Adopting the opinion of Dr Chapman, I should say Truth lies, in this controversy, nearly between the extremes. Then a moderate degree of nausea will best ensure the proper effect, though ~~this~~ it may sometimes be attained when no action is observable on the stomach. ^{H.}

As respects the other diuretics almost the whole might be employed in dropsy, but usage seems to have consecrated only a small number for its management, though perhaps the whole on some occasions may be brought into use.

The Digitalis holds a high station in the list of remedies against dropsy. It is one of the most powerful diuretics, and requires a cautious administration, as it sometimes produces effects highly unpleasant and alarming. The Digitalis is now allowed to be best adapted to those cases in which the force of arterial action has been taken off

L. Chapman's Therap. & Mat. Med. Vol. 1st p. 250.

by pouring
given in suc-
infusion or
the dose for-
ing and nig-
the infusion
a day. The
out drops
taken in the
according to S
dose is five
and London
half the do-
given in do-
or the same

I have also
cytalis requires
available effe-
days, when so-
alarming cha-
of strength, &
then circum-
cases to begin
the quantity of
is closely wat-

by previous depletion^{Le}. The dried leaves may be given in substance, or they may be taken in infusion or tincture. When taken in substance the dose for an adult is one grain morning and night. One table spoonful of the infusion may be given three times a day. The tincture being made of different degrees of strength is not uniformly taken in the same quantity. When made according to Garvins directions the medium dose is five drops. That of the Edinburgh and London Pharmacopoeias, possesses but half the strength, and is consequently given in double the quantity of Garvins or the saturated tincture.

I have already said the exhibition of digitalis requires circumspection. Sometimes no sensible effect follows its use for several days, when suddenly symptoms of the most alarming character arise, as extreme prostration of strength, great reduction of pulse &c. These circumstances make it necessary in all cases to begin with small doses increasing the quantity gradually, whilst the effect is closely watched until the object desired

Dr. Chapman's Treatise: & Mat. Med. Vol. 1st p. 245.

Dr. Ferriars Medical Histories p. 212.

be attained
during the
to void con
val days,
be discont
another s
the use of
puls, and
human f
changing
grated m
when the
ration, &c

The Su
medicine a
is imposed.
administrat
Ferriars, in
succeeded th
ced by his
rulers, to c
colled in
of Castor
in these
action.

be attained. Diuresis having been once induced, the patient will sometimes continue to void urine in large quantities for several days, even although the medicine be discontinued.

Another singular circumstance attending the use of digitalis is, that its effects on the pulse, and probably on other parts of the human frame may be made to vary by changing the posture of the body. The greatest reduction of pulse takes place when the person is in a recumbent position. &c.

The Super Saturated of Potash is another medicine in which the highest confidence is reposed. The success that followed its administration under the direction of Ferriar, in cases of dropsy generally, greatly exceeded that of any other article employed by him.th He directed from half an ounce, to an ounce and an half daily dissolved in ten ounces of water. The Oram of Castor is supposed to be most useful in those cases accompanied by febrile action.

The other
to, are the
Nitro, M.
Nematonic
substances
been men-
ed by diff-
substances
tions, and a
cepted to
-titution
themselves

In some
necessary to
ed, for as
inferior die
should be
perhaps has
practice a
some as an
Before
it is proper
was depends
be necessary

The other diuretics most frequently resorted to, are the carbonate of potash, *Spth Rth*: Nitrosi, Nitrate of potash Nistiana, Mela Nescatorius &c. These and all the preceding substances, with some others that have not been mentioned, are more or less recommended by different practitioners. Each of these substances enters into a variety of combinations, and in this way may be very usefully adapted to the different conditions of this constitution. ~~the constitution~~ that present themselves, as well as to the different periods of the disease.

In some cases of hydrothorax it becomes necessary to change the substance employed, for another of the same class, of known inferior diuretic powers. Why any advantage should be gained by such substitution, is perhaps hard to be explained; but in practice a knowledge of the fact often serves as an useful guide.

Before I close the remarks on diuretics it is proper to state, that when hydrothorax depends on visceral obstruction it will be necessary to remove this before a radi-

cal cure can
 in or kept up
 combined with
 at occasions
 productive of
 under such a
 often from the
 constitutes no
 With this to
 the cure of
 has been disc
 sivation of a
 ly inferior of
 as those ca
 depending on
 out of medicine
 There are also
 which, though
 requiring diet
 more debilitating
 arising under
 must be evid
 centiated by
 shew is to st
 The number of

cal cure can be made. It is in cases originating in, or kept up by obstruction, that mercury combined with diuretics and administered as at occasion a gentle ptyalism, has been productive of the most marked benefit.

Under such circumstances mercury alone will often prove highly beneficial, and sometimes completely remove the disease.

Hitherto the substances enumerated for the cure of hydrothorax, with one exception, have been diuretics. I now come to the consideration of another set of remedies of scarcely inferior efficacy;—There are *Tonics*.

As those cases of hydrothorax primarily depending on debility, however induced, this set of medicines is particularly adapted. There are also other instances of this disease, which, though at first inflammatory and requiring depletion, afterwards assume a more debilitated shape. Hydrothorax, existing under either of these forms, it must be evident can only be successfully combated by remedies whose general tendency is to strengthen the whole system.

The number of substances employed with

C. Maclean p. 126.

his intention
head of the
=parations,
=tables.

Maclean
of 1000 con
Myth in
to Griffith
exhibited in
from the
ation, that
diabetic an
that it is
arising fro
Under the
super is a
water with
enough to
flood the
ten to twenty
or three times
is given, the
much nausea
to open the
The vessel

this intention is comparatively small. At the head of them may be placed the chalybeate preparations, and next to these the tonic vegetables.

Maclean prefers as a tonic the sulphate of Iron combined with salt of tartar and Myrrh in the proportion recommended by Dr Griffiths. I have never seen this preparation exhibited in drops, but should suppose from the substances entering into its composition, that it is capable of combining a diuretic and tonic power, and therefore that it is particularly suited to dropsy arising from or connected with debility.

Under these circumstances the tonic I should prefer is a preparation formed by saturating water with sulphate of Iron and adding enough Sulphuric Acid to give to the fluid the acidity of Elixir Nitriol. From ten to twenty five drops may be given twice or three times a day. When the full dose is given, the stomach is occasionally so much nauseated that it becomes necessary to lessen the quantity.

The vegetable tonics are quina colom-

to the Throat
and the bit
sold in the
claries.

Of the
after

operation in
tried witho
suspension, it
form the op
thorax of
of tapping.
performed
after abrad
not much
where the op
fluid was
two other pa
Indeed, whe
contained in
be evacuated
by the proces
sification of the
stations &c. it
that the ope

bo &c. These singly, or variously combined, and the bitter and aromatic ingredients sold in the shops are all useful auxiliaries.

Of the Paracentesis Thoracis.

After all the medicines of known reputation in Hydrothorax have been fully tried without making any salutary impression, it has been recommended to perform the operation of paracentesis of the thorax, or of drawing off the water by means of tapping. This operation has been seldom performed in this country, and not very often abroad. The result of experience is not much in its favour. In those cases where the operation was performed, the fluid was found rapidly to reaccumulate, and the patient was finally carried off. Indeed, when the hydroptic fluid is contained in the pericardium, and cannot be evacuated by other means, or is supported by the presence of organic disease, as of inflammation of the vessels, aneurismal dilatation &c, it must appear evident, that the operation cannot be performed

with any rational hope of doing good.

London, 18th Dec 1801.

My Dear Sir,

Long quiet & all business gone home

Inaugural

Hyd

S. C. Kim

admitted Na